

Enhancement of the $K \rightarrow \pi$ Transition by the Instantaneous Weak Interactions

A.Z. Dubničková¹, S. Dubnička², V.N. Pervushin³, M. Sečanský^{3,2}

¹Department of Theoretical Physics, Comenius University, Slovak Republic

²Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovak Republic

³Joint Institute for Nuclear Research, 141980 Dubna, Russia

February 2, 2008

Abstract

The hadrodynamics of the instantaneous weak interactions (IWI) in Standard Model is considered supposing the resonance nature of the chiral hadronization of quark currents in QCD. Exploiting this supposition and QCD symmetries the IWI mechanism of enhancement of the $K \rightarrow \pi$ transition probability is obtained and relations between electroweak meson form factors and the amplitudes of the processes $K^+ \rightarrow \pi^+ e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$ and $K^+ \pi^- \rightarrow e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$ are established. We estimate possible parameters of the $K^+ \rightarrow \pi^+ e^+ e^-$ decay rate in NA48/2 CERN experiment using the chiral perturbation theory.

1. Introduction

The observation of kaon weak decays has been crucial for the modern theory of particle physics [1]. It was accepted to describe weak decays in the framework of electroweak theory at the quark QCD level including current vector boson weak interactions [2, 3]. However, a consistent theory at large distances of QCD is not yet constructed up to now. Therefore, now the most effective method of analysis of kaon decay physics [4, 5, 6, 7] is the chiral perturbation theory [8, 9]. In applying this method including one-loop contributions one should however take into account that the region of validity of the naive quantum chiral theory is only the low energy decomposition.

The first paper on the meson form factors in chiral perturbation theory [10] revealed the resonance nature of this chiral decomposition. Taking into account the summation of the chiral

series as the meson form factors in the investigation of the processes directly connected with the $K \rightarrow \pi$ transition (like nonleptonic kaon decays and $K^+ \rightarrow \pi^+ e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$) is the main difference of the effective chiral Lagrangian approach applied in the present paper from other ones cited above.

The introduction of the meson form factors in the chiral dynamics supposes that the low-energy dynamics can be separated by the Dirac formulation of the Standard Model [11, 12, 13, 14], where all vector fields (γ, W, Z) contain instantaneous potentials in the Lorentz frames defined by the *in* and *out* physical states considered in S-matrix as the irreducible representations of the Poincaré group. The dominance of the instantaneous interactions in the kaon-pion transitions differs our approach from the previous ones based on the heuristic Faddeev – Popov approach [15, 16], where instantaneous interactions were neglected.

We suppose also that the quark content of the mesons determines hadronization of QCD [17, 18] conserving its chiral and gauge symmetries.

These differences allow us to study the possibility of establishing connection of the corresponding decay probabilities with the π and K meson form factors and extracting information about form factors of π and K mesons from the $K^+ \rightarrow \pi^+$ transition in the $K^+ \rightarrow \pi^+ e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$ and $K^+ \pi^- \rightarrow e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$ processes. We present the explicit forms of the amplitudes of these processes in terms of meson form factors in the Standard Model of electroweak (EW) interactions.

2. Chiral Bosonization of EW Interaction

We begin with the Lagrangian of EW interaction of quarks

$$\mathcal{L}_{(J)} = -\frac{e}{2\sqrt{2}\sin\theta_W}(J_\mu^- W_\mu^+ + J_\mu^+ W_\mu^-), \quad (1)$$

where $J_\mu^+ = \bar{d}'\gamma_\mu(1 - \gamma_5)u$; $\bar{d}' = d\cos\theta_C + s\sin\theta_C$, θ_C is a Cabbibo angle $\sin\theta_C = 0.223$. The quark content of π^+ and K^+ mesons $\pi^+ = (\bar{d}, u)$, $K^+ = (\bar{s}, u)$, $\bar{K}^0 = (\bar{s}, d)$ leads to the effective chiral hadron currents $J_\mu^\pm$ in the Lagrangian (1)

$$J_\mu^\pm = [J_\mu^1 \pm iJ_\mu^2] \cos\theta_C + [J_\mu^4 \pm iJ_\mu^5] \sin\theta_C, \quad (2)$$

where using the Gell-Mann matrixes λ^k one can define the meson current as [8]

$$i \sum \lambda^k J_\mu^k = i\lambda^k (V_\mu^k - A_\mu^k)^k = F_\pi^2 e^{i\xi} \partial_\mu e^{-i\xi}, \quad (3)$$

$$\xi = F_\pi^{-1} \sum_{k=1}^8 M^k \lambda^k = F_\pi^{-1} \begin{pmatrix} \pi^0 + \frac{\eta}{\sqrt{3}} & \pi^+ \sqrt{2} & K^+ \sqrt{2} \\ \pi^- \sqrt{2} & -\pi^0 + \frac{\eta}{\sqrt{3}} & K^0 \sqrt{2} \\ K^- \sqrt{2} & \bar{K}^0 \sqrt{2} & -\frac{2\eta}{\sqrt{3}} \end{pmatrix}. \quad (4)$$

In the first orders in mesons one can write

$$V_\mu^- = \sqrt{2} [\sin\theta_C (K^- \partial_\mu \pi^0 - \pi^0 \partial_\mu K^-) + \cos\theta_C (\pi^- \partial_\mu \pi^0 - \pi^0 \partial_\mu \pi^-)] \quad (5)$$

and

$$A_\mu^- = \sqrt{2} F_\pi (\partial^\mu K^- \sin\theta_C + \partial^\mu \pi^- \cos\theta_C); \quad (6)$$

here $F_\pi \simeq 92$ MeV. The right form of chiral Lagrangian of the electromagnetic interaction of mesons can be constructed by the covariant derivative $\partial_\mu \chi^\pm \rightarrow D_\mu \chi^\pm \equiv (\partial_\mu \pm ieA_\mu) \chi^\pm$, where $\chi^\pm = K^\pm, \pi^\pm$.

The main idea is a consideration of the instantaneous interactions in this model (that is associated with the Yukawa potential of massive vector bosons).

3. Instantaneous Coulomb Interaction in QED

In order to explain the main idea of describing of massive bosons, let us consider the action of photons $S_{\text{QED}} = \int d^4x \mathcal{L}$, where

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F_{\mu\nu} + j_\mu A_\mu - \bar{\psi}(i\gamma\partial + m)\psi \quad (\gamma\partial = \gamma_\mu \partial_\mu), \quad (7)$$

is the QED Lagrangian, A_μ is a vector potential, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, index $\mu, \nu = 0, 1, 2, 3$, ψ is the electron-positron field described by the Dirac bispinor, $j_\mu = e\bar{\psi}\gamma_\mu\psi$ is the charge current, and e is the electron charge.

The instantaneous Coulomb interaction is an attribute of the Hamiltonian formulation in a concrete Lorentz frame in QED, because the time component of the vector field A_0 has the zero canonical momentum¹ $P_0 = \partial\mathcal{L}/\partial(\partial_0 A_0) = 0$. Therefore, Dirac [11] supposed to consider A_0 as a classical field and excluded it by the manifest resolution of the Gauss constraint $\Delta A_0 - \partial_0 \partial_k A_k = -j_0$ (where $\Delta = \partial_j^2$) and the gauge transformation

$$A_0^{(\text{rad})}(A) = A_0 + \partial_0 \lambda^{(\text{rad})}(A) = A_0 - \partial_0 \frac{1}{\Delta} \partial_k A_k, \quad (8)$$

$$A_l^{(\text{rad})}(A) = A_l + \partial_l \lambda^{(\text{rad})}(A) = A_l - \partial_l \frac{1}{\Delta} \partial_k A_k, \quad (9)$$

$$\psi^{(\text{rad})}(\psi, A) = e^{ze\lambda^{(\text{rad})}(A)} \psi, \quad (10)$$

where $\lambda^{(\text{rad})}(A) = -\frac{1}{\Delta} \partial_k A_k \equiv \frac{1}{4\pi} \int d^3y \frac{\partial_k A_k(x_0, y_k)}{|x-y|}$. These gauge-invariant functionals were called the “dressed fields”, or the “radiation variables”.

In terms of the gauge-invariant radiation variables the initial action of QED takes a form

$$S_{\text{QED}}^{(\text{rad})} = \int d^4x \left[\frac{1}{2} (\partial_\mu A_k^{(\text{rad})})^2 + \frac{1}{2} j_0^{(\text{rad})} \frac{1}{\Delta} j_0^{(\text{rad})} - A_k^{(\text{rad})} j_k^{(\text{rad})} - \bar{\psi}^{(\text{rad})} (i\gamma\partial + M) \psi^{(\text{rad})} \right]. \quad (11)$$

This Lagrangian contains the Coulomb instantaneous interaction forming atomic and molecular bound states in the lowest order of the “radiation corrections” $A_k^{(\text{rad})} = 0$. The radiation action (11) corresponds to the propagator

$$D_{\mu\nu}^{\gamma,(\text{rad})}(q) = \delta_{\mu 0} \delta_{\nu 0} \frac{1}{\vec{q}^2} + \delta_{\mu i} \delta_{\nu j} \left(\delta_{ij} - \frac{q_i q_j}{\vec{q}^2} \right) \frac{1}{q^2 + i\varepsilon} \equiv \quad (12)$$

$$\equiv -\frac{\delta_{\mu\nu}}{q^2 + i\varepsilon} + \frac{(q_0 \delta_{\mu 0})(q_0 \delta_{\nu 0}) - (q_i \delta_{\mu i})(q_j \delta_{\nu j})}{\vec{q}^2 [q^2 + i\varepsilon]}. \quad (13)$$

Recall that Faddeev [16] considered this Dirac approach (11) to QED applied by Schwinger [19] for non-Abelian theory as the foundation of the Faddeev – Popov (FP) heuristic approach [15],

¹Here $A_0 = A_\mu n_\mu$ and n_μ is a unit timelike vector $n_\mu^2 = 1$ that distinguishes timelike and spacelike components $[A_\mu]_n = [(A_\mu n_\mu), A_\nu - n_\nu (A_\mu n_\mu)]_{n_\mu=(1,0,0,0)} = [A_0, A_k]$.

where the instantaneous interactions disappear. Really, one can prove [16] that the last (longitudinal) term in Eq. (13) can be neglected for the class of the elementary particle scattering amplitudes, where charge currents are conserving and satisfy the equality $j_0 q_0 - j_k q_k = 0$. However, in the case of calculation of the bound state spectrum the last term cannot be neglected, because it gives in (12) the instantaneous singularity forming observable atoms.

Using the Lorentz transformations (proved at the level of Poincare algebra of observables [20]) one can find the photon propagator in an arbitrary frame as follows [12],

$$D_{\mu\nu}^{\gamma,(\text{rad})}(q|n) = \frac{n_\mu n_\nu}{|q^\perp|^2} - \left(\delta_{\mu\nu}^\perp - \frac{q_\mu^\perp q_\nu^\perp}{|q^\perp|^2} \right) \frac{1}{q^2 + i\epsilon}, \quad (14)$$

where $q_\mu^\perp = q_\mu - n_\mu(qn)$, $\delta_{\mu\nu}^\perp = \delta_{\mu\nu} - n_\mu n_\nu$, and n_μ is the timelike vector reflecting the initial conditions defined by physical states of the quantized fields [14, 17, 18].

4. Instantaneous Interaction in SM

SM was formulated [21] on the basis of the FP heuristic approach [15] without instantaneous interactions. Therefore, we shall use here the generalization of the fundamental radiation variables in QED (14) to massive vector fields, described by the action,

$$W = \int d^4x \left[-\frac{1}{2}(\partial_\mu W_\nu^+ - \partial_\nu W_\mu^+)(\partial_\mu W_\nu^- - \partial_\nu W_\mu^-) + M_W^2 W_\mu^+ W_\mu^- + J_\mu^- W_\mu^+ + J_\mu^+ W_\mu^- \right]$$

that considered in [13] using the manifest resolution of the Gauss equation and “dressing” all fields by the gauge transformation with $\lambda^{\pm(\text{rad})} = [1/(M_W^2 - \Delta)]\partial_k W_k^\pm$.

The action of W-bosons in SM in terms of the radiation variables contains the instantaneous Yukawa interaction of the currents (5) and (6) with the Hamiltonian

$$H_{\text{Yuk}} = \frac{G_F}{\sqrt{2}} \int d^3x J_0^- \frac{1}{M_W^2 - \Delta} J_0^+, \quad \text{where} \quad \frac{G_F}{\sqrt{2}} = \frac{e^2}{8 \sin^2 \theta_W}. \quad (15)$$

The radiation variables lead to a propagator of type of the QED one (12)

$$D_{\mu\nu}^{W,(\text{rad})}(q) = \delta_{\mu 0} \delta_{\nu 0} \frac{1}{q^2 + M_W^2} + \delta_{\mu i} \delta_{\nu j} \left(\delta_{ij} - \frac{q_i q_j}{q^2 + M_W^2} \right) \frac{1}{q^2 - M_W^2 + i\epsilon}. \quad (16)$$

which has no singularity in the massless limit and is well behaved for large momenta. As it was shown in [13], the Lorentz transformations of classical radiation variables coincide with the quantum ones and they both (quantum and classical) correspond to the transition to another Lorentz reference frame distinguished by another time-axis, where the propagator takes the form

$$D_{\mu\nu}^{W,(\text{rad})}(q|n) = \frac{n_\mu n_\nu}{M_W^2 + |q^\perp|^2} - \left(\delta_{\mu\nu}^\perp - \frac{q_\mu^\perp q_\nu^\perp}{M_W^2 + |q^\perp|^2} \right) \frac{1}{q^2 - M_W^2 + i\epsilon} \quad (17)$$

with $q_\mu^\perp = q_\mu - n_\mu(qn)$, $\delta_{\mu\nu}^\perp = \delta_{\mu\nu} - n_\mu n_\nu$.

5. The $K \rightarrow \pi$ Transition Amplitude and the Rule $\Delta T = \frac{1}{2}$

One can estimate the $K^+ \rightarrow \pi^+$ transition amplitude

$$\langle \pi^+ | -i \int dx^0 H_{\text{Yuk}} | K^+ \rangle = i(2\pi)^4 \delta^4(k-p) G_{\text{EW}} \Sigma(p^2), \quad (18)$$

where

$$G_{\text{EW}} = \frac{\sin \theta_C \cos \theta_C}{8M_W^2} \frac{e^2}{\sin^2 \theta_W} \equiv \sin \theta_C \cos \theta_C \frac{G_F}{\sqrt{2}} \quad (19)$$

is the constant. Making the normal ordering of the π^0 fields

$$\Delta(\vec{x} - \vec{y}) = \langle 0 | \pi^i(x) \pi^{i'}(y) | 0 \rangle = \delta^{ii'} \int \frac{d^3 l}{(2\pi)^3} \frac{e^{i\vec{l} \cdot (\vec{x} - \vec{y})}}{2E_\pi(\vec{l})}, \quad (20)$$

where $E_\pi(\vec{l}) = \sqrt{m_\pi^2 + \vec{l}^2}$ is the energy of π -meson, in the product of vector currents, one can find the expression in the lowest order of chiral perturbation theory

$$\Sigma(p^2) = 2p_0^2 \left[\frac{F_\pi^2}{M_W^2} + \frac{1}{(2\pi)^3} \int \frac{d^3 l}{2E_\pi(\vec{l})} \frac{1}{M_W^2 + \vec{l}^2} \right] \equiv 2p_0^2 \frac{F_\pi^2}{M_W^2} g_8; \quad (21)$$

here

$$g_8 = 1 + \frac{M_W^2}{F_\pi^2 (2\pi)^3} \int \frac{d^3 l}{2E_\pi(\vec{l})} \frac{1}{M_W^2 + \vec{l}^2} \quad (22)$$

is the parameter of the enhancement of the probability of $K^+ \rightarrow \pi^+$ transition. In the low-energy chiral perturbation theory [10], the summation of the chiral series can be parameterized by the meson form factors $f_{K\pi W}^V(-\vec{l}^2)$ and $f_{\pi\pi W}^V(-\vec{l}^2)$ [4, 6, 7].

Finally, the instantaneous interaction can give the result

$$g_8 = 1 + \frac{M_W^2}{F_\pi^2 (2\pi)^3} \int \frac{d^3 l}{2E_\pi(\vec{l})} \frac{f_{K\pi W}^V(-\vec{l}^2) f_{\pi\pi W}^V(-\vec{l}^2)}{M_W^2 + \vec{l}^2}, \quad (23)$$

where $g_8 = 5.1$ [5]. The relation (23), can be considered as the low-energy sum rule for meson form factors in the space-like region.

This result shows that the simultaneous vector interaction can increase the axial interaction $K^+ \rightarrow \pi^+$ transition in g_8 times and give a new term describing the $K^0 \rightarrow \pi^0$ transition proportional to $g_8 - 1$.

Using the covariant perturbation theory [22]² and relativistic invariance ($J_0^2 \rightarrow J_\mu^2$) one can calculate this enhancement in terms of currents as the effective Lagrangians [23]

$$\mathcal{L}_{(\Delta T = \frac{1}{2})} = \frac{G_F}{\sqrt{2}} g_8 \cos \theta_C \sin \theta_C \left[(J_\mu^1 + iJ_\mu^2)(J_\mu^4 - iJ_\mu^5) - (J_\mu^3 + \frac{1}{\sqrt{3}}J_\mu^8)(J_\mu^6 - iJ_\mu^7) + h.c. \right], \quad (24)$$

$$\mathcal{L}_{(\Delta T = \frac{3}{2})} = \frac{G_F}{\sqrt{2}} \cos \theta_C \sin \theta_C \left[(J_\mu^3 + \frac{1}{\sqrt{3}}J_\mu^8)(J_\mu^6 - iJ_\mu^7) + h.c. \right] \quad (25)$$

in satisfactory agreement with experimental data within the accuracy $20 \div 30\%$ [8, 23].

²The covariant perturbation theory was developed as the series $J_0^k(\gamma \oplus \xi) = J_0^k(\xi) + F_\pi^2 \partial_0 \gamma^k + \gamma^i f_{ijk} J_0^j(\xi) + O(\gamma^2)$ with respect to quantum fields γ added to ξ as the product $e^{i\gamma} e^{i\xi} \equiv e^{i(\gamma \oplus \xi)}$ [22].




Fig. 1: Axial current contribution

6. The $K^+ \rightarrow \pi^+ + l + \bar{l}$ amplitude

The result of calculation of the axial contributions to the amplitude of the process $K^+ \rightarrow \pi^+ + l + \bar{l}$ within the framework of chiral Lagrangian (1), (3) including phenomenological meson form factors shown in Fig. 1 as a fat dot takes the form

$$T^{AA}(K^+ \rightarrow \pi + l^+ l^-) = 2eG_{EW}L_\nu D_{\mu\nu}^{\gamma(rad)}(q)(k_\mu + p_\mu) t^{AA}(q^2, k^2, p^2), \quad (26)$$

where G_{EW} is the constant (19), $L_\mu = \bar{l}\gamma_\mu l$ is leptonic current,

$$t^{AA}(q^2, k^2, p^2) = F_\pi^2 \left[\frac{f_\pi^V(q^2)k^2}{m_\pi^2 - k^2 - i\epsilon} + \frac{f_K^V(q^2)p^2}{M_K^2 - p^2 - i\epsilon} + \frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} \right], \quad (27)$$

and $f_{\pi,K}^{(A,V)}(q^2)$ are meson form factors. In this expression the propagator $D_{\mu\nu}^{W,(rad)}(p|n)$ given by Eq. (17) keeps only the Yukawa potential part $\frac{p_\mu p_\nu}{p^2 M_W^2}$.

On the mass-shell the sum (27) takes the form

$$\begin{aligned} t^{AA}(q^2, M_K^2, m_\pi^2) &= \\ &= t(q^2) = F_\pi^2 \left[\frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right]. \end{aligned} \quad (28)$$

As can be seen in [4, 5], the amplitude for $K^+ \rightarrow \pi^+ + l + \bar{l}$ vanishes at the tree level, where form factors are equal to unit $t(q^2)|_{f^V=f^A=1} = 0$. It is well known that these form factors in the limit $q^2 \rightarrow 0$ take the form $f_{\pi,K}^{A,V}(q^2) = 1 - \frac{q^2}{6} < r^2 >_{\pi,K}^{(A,V)}$, where $< r^2 >_{\pi,K}^{(A,V)}$ are the axial and vector mean square radii of mesons, respectively, in agreement with the chiral perturbation theory [4, 8, 10].

The result of calculation of vector contributions to the amplitude of the process $K^+ \rightarrow \pi^+ + l + \bar{l}$ within the framework of chiral Lagrangian (1), (3) including phenomenological meson form factors shown in Fig. 2 as a fat dot takes the form

$$T^{VV(rad)}(K^+ \rightarrow \pi + l^+ l^-) = 2eG_{EW}L_\nu D_{\mu\nu}^{\gamma(rad)}(q)(k_\mu + p_\mu) t^{VV}(q^2, k^2, p^2), \quad (29)$$




Fig. 2: Vector current contribution

where

$$t^{VV}(q^2, k^2, p^2) = \frac{1}{2(2\pi)^2} \int \frac{d|\vec{l}|}{E_\pi(\vec{l})} \frac{\vec{l}^2}{E_\pi(\vec{l})} \left\{ f_{K\pi W}^V(\vec{l}^2) f_{\pi\pi W}^V(\vec{l}^2) \left[\frac{f_\pi^V(q^2) k^2}{m_\pi^2 - k^2 - i\epsilon} + \frac{f_K^V(q^2) p^2}{M_K^2 - p^2 - i\epsilon} \right] + \frac{f_\pi^A(q^2) f_{K\pi W}^V(\vec{l}^2) f_{\pi\pi W\gamma}^V(\vec{l}^2) + f_K^A(q^2) f_{\pi\pi W}^V(\vec{l}^2) f_{K\pi W\gamma}^V(\vec{l}^2)}{2} \right\};$$

here $f_{\pi\pi W\gamma}^V(\vec{l}^2)$, $f_{K\pi W\gamma}^V(\vec{l}^2)$ are the four particle interaction form factors which should coincide with the three particle interaction form factors $f_{\pi\pi W}^V = f_{\pi\pi W}^V$, $f_{K\pi W\gamma}^V = f_{K\pi W}^V$ in accordance with the gauge invariance of the strong interactions.

Using the result (22) one can write on the mass shell $k^2 = M_K^2, p^2 = m_\pi^2$:

$$t^{VV}(q^2, M_K^2, m_\pi^2) = (g_8 - 1)t(q^2), \quad (30)$$

$$t^{AA}(q^2, M_K^2, m_\pi^2) + t^{VV}(q^2, M_K^2, m_\pi^2) = g_8 t(q^2), \quad (31)$$

where $g_8, t(q^2)$ are given by Eqs. (22) and (28), respectively.

Finally, the result of calculation of axial and vector contributions to the amplitude of the process $K^+ \rightarrow \pi^+ + l + \bar{l}$ within the framework of chiral Lagrangian (2),(5) including phenomenological meson form factors takes the form

$$T_{(K^+ \rightarrow \pi^+ l^- \bar{l}^-)}^{(\text{rad})} = 2eg_8 G_{EW} L_\nu D_{\mu\nu}^{\gamma(\text{rad})}(q)(k_\mu + p_\mu) t(q^2). \quad (32)$$

This amplitude leads to the total decay rate for the transition $K^+ \rightarrow \pi^+ e^+ e^-$

$$\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)}((M_K - m_\pi)^2) = \bar{\Gamma}_{e^+ e^-} \int_{4m_e^2}^{(M_K - m_\pi)^2} \frac{dq^2}{M_K^2} \rho(q^2) F(q^2), \quad (33)$$

where

$$\bar{\Gamma}_{e^+ e^-} = \frac{(s_1 c_1 c_3)^2 g_8^2 G_F^2 \alpha^2 M_K^5}{(4\pi)^4 24\pi} \Big|_{g_8=5.1} = 1.37 \times 10^{-22} \text{GeV}, \quad (34)$$

$$\begin{aligned}
F(q^2) &= \left[\frac{(4\pi)^2 t(q^2)}{q^2} \right]^2 = \\
&= \left[\frac{(4\pi F_\pi)^2}{q^2} \right]^2 \left[\frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right]^2 \quad (35)
\end{aligned}$$

and

$$\rho(q^2) = \left(1 - \frac{4m_l^2}{q^2} \right)^{1/2} \times \left(1 + \frac{2m_l^2}{q^2} \right) \lambda^{3/2}(1, q^2/M_K^2, m_\pi^2/M_K^2);$$

here $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + bc + ca)$, $s_1 c_1 c_3$ is the product of Cabibbo-Kobayashi-Maskawa matrix elements $V_{ud}V_{us}$.

The mechanism of enhancement $|\Delta T| = \frac{1}{2}$ considered in this paper can be generalized to a description of other processes including $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ by replacing the γ -propagator by the Z -boson one, so that we have the relations

$$\begin{aligned}
&\left[\frac{q^2}{(4\pi F_\pi)^2} \right]^2 \frac{M_K^2}{\rho(q^2) \bar{\Gamma}_{e^+e^-}} \frac{d\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)}(q^2)}{dq^2} = \\
&= \left[\frac{M_Z^2}{(4\pi F_\pi)^2} \right]^2 \frac{M_K^2}{\rho(q^2) \bar{\Gamma}_{\nu\bar{\nu}}} \frac{d\Gamma_{(K^+ \rightarrow \pi^+ \nu \bar{\nu})}(q^2)}{dq^2} = \\
&= \left[\frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right]^2. \quad (36)
\end{aligned}$$

Thus, exploiting the instantaneous weak interaction mechanism of enhancement in the $K \rightarrow \pi$ transition probability and QCD symmetry we derive the sum rule of EW vector meson form factors given by Eqs. (22), (28), (33) and their relation to the differential $K \rightarrow \pi e^+ e^-$ decay rate (36).

6.1. The form factor probe of the differential $K \rightarrow \pi e^+ e^-$ decay rate

The estimation of the meson loop contribution was made in [5] where a function $\hat{\phi}^2(q^2)$ was used instead of the form factor rate $F(q^2)$ (35)

$$F(q^2) \implies \hat{\phi}^2(q^2)$$

It was shown that the values $g_8 = 5.1$, $\hat{\phi}(0) = 1$ gave the total decay rate $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.91 \times 10^{-23}$ GeV in the satisfactory agreement with the experimental data $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.44 \pm 0.27 \times 10^{-23}$ GeV [24, 25]. However, the main contribution goes from the baryon loops [4, 6, 7]. Therefore, we discuss here the value of the differential $K \rightarrow \pi e^+ e^-$ decay rate (35) in the chiral perturbation theory [8, 10] where both the pion loop contribution Π_π and baryon ones lead to the meson form factors and resonances [4, 10] in the Padé-type approximation

$$f_\pi^V(q^2) = 1 + M_\rho^{-2} q^2 + \alpha_0 \Pi_\pi(q^2) + \dots \simeq \frac{1}{1 - M_\rho^{-2} q^2 - \alpha_0 \Pi_\pi(q^2)} \quad (37)$$

$$f_\pi^A(q^2) \simeq f_K^A(q^2) = 1 + M_a^{-2} q^2 + \dots \simeq \frac{1}{1 - M_a^{-2} q^2} \quad (38)$$

where

$$\alpha_0 = \frac{m_\pi^2}{(4\pi F_\pi)^2} \frac{4}{3} \simeq \frac{2}{103}, \quad (39)$$

$$\Pi_\pi(t) = (1 - \bar{t}) \left(\frac{1}{\bar{t}} - 1 \right)^{1/2} \arctan \left(\frac{\bar{t}^{1/2}}{(1 - \bar{t})^{1/2}} \right) - 1; \quad 0 < t < (2m_\pi)^2 \quad (40)$$

$$\Pi_\pi(t) = \frac{\bar{t} - 1}{2} \left(1 - \frac{1}{\bar{t}} \right)^{1/2} \left\{ i\pi - \log \frac{\bar{t}^{1/2} + (\bar{t} - 1)^{1/2}}{\bar{t}^{1/2} - (\bar{t} - 1)^{1/2}} \right\} + 1; \quad (2m_\pi)^2 < t \quad (41)$$

is the pion loop contribution [8, 10], and $M_\rho = 771$ MeV and $M_a = 980$ MeV are the values of resonance masses [24] that can be determined by the baryon loops [4, 6, 7, 10].

The constant approximation of the type of [5]

$$F(q^2 = 0) = F(0) = (4\pi F_\pi)^4 \frac{[f_\pi^A(q^2) - f_\pi^V(q^2)]^2}{(q^2)^2} \Big|_{q^2=0} = (4\pi F_\pi)^4 [M_\rho^{-2} - M_a^{-2}]^2 = 0.74 \quad (42)$$

corresponds to the value of $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.41 \times 10^{-23}$ GeV in satisfactory agreement with the experimental data $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.44 \pm 0.27 \times 10^{-23}$ GeV [24, 25].

The form factors (37) and (38) below the two particle threshold $q^2 < 4m_\pi^2$ determines the differential rate $F(q^2)$ as a function of q^2

$$F(q^2) = \frac{F(0)}{[1 - M_\rho^{-2} q^2]^2 [1 - M_a^{-2} q^2]^2} \longrightarrow \frac{d}{dq^2} \log F(q^2) \Big|_{q^2=0} \simeq 2[M_\rho^{-2} + M_a^{-2}] \simeq 5 \text{ GeV}^{-2} \quad (43)$$

At the region above the two particle threshold $4m_\pi^2 < q^2 < (M_K - m_\pi)^2$ there is a jump of the differential rate $F(q^2)$ at the level of $5 \div 10$ %

$$F(q^2) = (4\pi F_\pi)^4 \frac{[M_\rho^{-2} - M_a^{-2}]^2 + \alpha_0^2 \pi^2 (q^2 - 4m_\pi^2)/(4q^6)}{[1 - M_\rho^{-2} q^2]^2 [1 - M_a^{-2} q^2]^2} \quad (44)$$

These results (42), (43), and (44) are arguments that the detailed investigation of the differential $K \rightarrow \pi e^+ e^-$ decay rate in the NA48/2 CERN experiment can give us information about the meson form factors. To make much more realistic analysis, we have to use the unitary and analytic model for meson form factors [26] describing very well all experimental data.

Conclusion

We have seen that experimental data testify to that the kaon-pion transition can be obtained, in SM model, by the normal ordering of the pion field in the instantaneous interaction (15) as one of the indispensable attributes of the Dirac approach to SM [14]. This interaction can explain the rule $\Delta T = 1/2$ in the nonleptonic kaon decays in the lowest order of the “radiation corrections”. Paraphrasing Lord Eddington: “A proton yesterday and electron today do not make an atom” [27] one can say that “A kaon yesterday and pion today do not make a weak $K \rightarrow \pi$ transition” [2, 3]. The dominance of the instantaneous interaction of massive vector boson time components justifies the application of the chiral perturbation theory leading to meson form factors [8, 10]. Exploiting the instantaneous weak interaction (IWI) mechanism of enhancement in the $K \rightarrow \pi$ transition probability and QCD symmetry we derive the relation between meson form factors and the differential semileptonic decay rates (36). These relations allow us to estimate parameters of $K \rightarrow \pi e^+ e^-$ decay rate in the NA48/2 CERN experiment.

These results show us that the Dirac Hamiltonian formulation [13] of Standard Model can reveal new physical effects in the comparison with its FP heuristic version [21] used now for the describing observational data. Therefore, the problem of such the formulation becomes topical in the light of the future precision experiments.

Acknowledgments

The authors are grateful to B.M. Barbashov, D.Yu. Bardin, A.Di Giacomo, S.B. Gerasimov, G.V. Efimov, A.V. Efremov, V.D. Kekelidze, E.A. Kuraev, A.N. Sissakian and M.K. Volkov for fruitful discussions. The work was in part supported by the Slovak Grant Agency for Sciences VEGA, Gr.No.2/4099/26.

References

- [1] I. Mikulec, Kaon Physics, XXIV Physics in Collision - Boston, June 27-29, 2004.
- [2] G. Altarelli and L. Maiani, Phys. Lett. **52B**, 351 (1974).
M. K. Gaillard and B. W. Lee, Phys. Rev. Lett. **33**, 108 (1974).
- [3] A. I. Vainshtein et al., Yad. Fiz. **24**, 820 (1976);
S. S. Gershtein and M. Yu. Khlopov, JETP Lett., **23**, 338 (1976).
- [4] A. A. Belkov, Yu. L. Kalinovsky, V. N. Pervushin, Description of $K^+ \rightarrow \pi^+ e^+ e^-$ Decay in Quantum Chiral Theory. JINR-P2-85-107, Feb. 1985. (Dubna, JINR);
A. A. Bel'kov et al., Sov. J. Nucl. Phys. **44**, 690 (1986).
- [5] G. Ecker et al., Nucl. Phys. **B 291**, 692 (1987).
- [6] A. A. Bel'kov et al., Phys. Part. Nucl. **26**, 239 (1995).
- [7] A. A. Bel'kov et al., Phys. Lett. **220 B**, 459 (1989).
- [8] M. K. Volkov and V. N. Pervushin, Usp. Fiz. Nauk **120**, 363 (1976);
M. K. Volkov and V. N. Pervushin, "Essentially Nonlinear Field Theory, Dynamical Symmetry and Pion Physics", Atomizdat, Moscow, ed. D. I. Blokhintsev, 1979 (in Russian).
- [9] J. Gasser and H. Leutwyler, Ann. of Phys. **158**, 142 (1984).
- [10] M. Volkov, V. Pervushin, Phys. Lett. **51B**, 356 (1974).
- [11] P. A. M. Dirac, Proc. Roy. Soc., **A 114**, 243 (1927);
P. A. M. Dirac, Can. J. Phys., **33**, 650 (1955).
- [12] I. V. Polubarinov, Physics of Particles and Nuclei, **34**, 377 (2003).
- [13] H.-P. Pavel and V. N. Pervushin, Int. J. Mod. Phys. **A 14**, 2885 (1999).
- [14] A. Z. Dubničková, S. Dubnička, V.N. Pervushin, M. Sečansky, hep-ph/0607211.
- [15] L. Faddeev and V. Popov, Phys. Lett. **25 B**, 29 (1967).
- [16] L. Faddeev, T. M. F., **1**, 3 (1969).
- [17] Yu. L. Kalinovsky et al., Sov. J. Nucl. Phys., **49**, 1709 (1989);
V. N. Pervushin, Nucl. Phys. B (Proc. Supp.) **15**, 197 (1990).
- [18] V. N. Pervushin, Physics of Particles and Nuclei, **34**, 348 (2003).

- [19] J. Schwinger, Phys. Rev., **127**, 324 (1962).
- [20] B. Zumino, J. Math. Phys., **1**, 1 (1960).
- [21] D. Bardin and G. Passarino, “The standard model in the making: precision study of the electroweak interactions”, Clarendon, Oxford, 1999.
- [22] V. N. Pervushin, Theor. Math. Phys., **27**, 330 (1977).
- [23] Yu. L. Kalinovsky, V. N. Pervushin, Sov. J. Nucl. Phys., **29**, 475 (1979).
- [24] Review of Particle Physics, Phys. Lett., **592 B**, 1 (2004).
- [25] P. Bloch et al., Phys. Lett., **56 B**, 201 (1975).
- [26] M. E. Biagini, S. Dubnička, E. Etim, P. Kolar, Nuovo Cim., **A 104**, 363 (1991);
S. Dubnička, A. Z. Dubničková, A. Antušek, Acta Physica Slovaca, **49**, 165 (1999);
S. Dubnička, A. Z. Dubničková, J. Phys., **G28**, 2137 (2002).
- [27] A. Eddington, “Relativistic Theory of Protons and Electrons“, 1936, Cambridge University Press.